

**PRESCRIBER'S ORDERS
FOR EMERGENCY DEPARTMENT
PROCEDURAL SEDATION / ANALGESIA**

DATE / /
DD MM YYYY

TIME : HOURS
HH MM

WEIGHT kilograms HEIGHT centimetres ALLERGY CAUTION sheet reviewed

Pharmacy
Use Only

WRITE FIRMLY WITH A BALLPOINT PEN WITH BLUE OR BLACK INK

Noted by
RN/UC

1. Oral (PO) Sedation : administer 20-45 minutes prior to procedure
 Midazolam mg PO x 1 dose (0.5-1 mg/kg/dose. Maximum 20 mg)

2. Intravenous (IV) Sedation
 Ketamine mg IV x 1 dose (1 mg/kg/dose. Maximum 75 mg)
2nd dose:
 Ketamine mg IV x 1 dose (0.5-1 mg/kg/dose. Maximum 75 mg)

OR

Midazolam mg IV x 1 dose (0.1-0.2 mg/kg/dose. Maximum 8 mg)

Fentanyl micrograms IV, may be repeated every minutes for total of 3 doses. (1 microgram /kg/dose. Maximum 150 micrograms)

Other: _____

3. Adjunct Drugs: calculate appropriate doses
Drugs to be available only, NOT to be drawn up unless needed.

- A) Atropine mg IV. (0.01 mg/kg/dose. Maximum 0.5 mg)
- B) Flumazenil mg IV. (0.01 mg/kg/dose. Maximum 1 mg)
- C) Naloxone mg IV. (0.01 mg/kg/dose. Maximum 2 mg)
- D) Ondansetron mg IV. (0.15 mg/kg/dose. Maximum 8 mg)
- E) Succinylcholine mg IV. (1 mg/kg/dose)

4. Monitoring guidelines:

- RR and depth until unstimulated Arousal Score 2 or less by dedicated person
- Continuous SpO₂ and heart rate monitoring
- Record observations (Heart rate, RR, SpO₂, Arousal Score) on sedation/analgesia record following sedation and during recovery at least every 5 minutes until the pre-procedure level of consciousness is resumed and then every 15 minutes until the Arousal Score is 2 or less.
- See BCCH Sedation/ Analgesia Guidelines and form CH 436

All dose formulations, calculations and preparations to be independently double checked by 2 clinicians (nurses and/or physicians).

Signature: _____ Pager # _____

Print Name: _____ College ID# _____