



†Pre-discharge imaging indicated if contrast blush or evidence of psuedoaneurysm on initial CT

ΩNo post discharge imaging if patient asymptomatic and prior imaging showed no evidence of psuedoaneurysm

¥Return to full impact activities at the discretion of the managing surgeon

References

- Holevar M, DiGiacomo JC, Ebert J, et al. Practice management guidelines for the management of genitourinary trauma. Eastern Association for the Surgery of Trauma. 2004. <http://www.east.org/resources/treatment-guidelines/genitourinary-trauma-management-of>
- Holmes JF, Sokolove PE, Brant WE, et al. Identification of children with intra-abdominal injuries after blunt trauma. Ann Emerg Med. 2002 May;39(5):500-9.
- Lindberg D, Makoroff K, Harper N, et al. Utility of hepatic transaminases to recognize abuse in children. Pediatrics. 2009 Aug;124(2):509-16.
- Karaduman D, Sarioglu-Buke A, Kilic I, Gurses E. The role of elevated liver transaminase levels in children with blunt abdominal trauma. Injury. 2003 May; 34(4):249-52.
- Karam O, La Scala G, Le Coultre C, Chardot C. Liver function tests in children with blunt abdominal traumas. Eur J Pediatr Surg. 2007 Oct;17(5):313-6.
- Puranik SR, Hayes JS, Long J, Mata M. Liver enzymes as predictors of liver damage due to blunt abdominal trauma in children. South Med J. 2002 Feb;95(2):203-6.
- Pediatric trauma society. Liver/spleen guideline examples.
<http://pediatrictraumasociety.org/benchmark/clinical-resources.cgi>
- Safavi A, Beudry P, Jamieson D, Murphy J. Traumatic pseudoaneurysms of the liver and spleen in children: is routine screening warranted? J Pediatr Surg. 2011 May; 46(5):938-41.